

reduced into Parallelism and diverge no more. For then those rays will recompose a Beam of white Light XY. If the refracting Angle of either Prism be the bigger, that Prism must be so much the nearer to the Lens. You will know when the Prisms and the Lens are well set together by observing if the Beam of Light XY which comes out of the second Prism be perfectly white to the very edges of the Light, and at all distances from the Prism continue perfectly and totally white like a Beam of the Sun's Light. For till this happens, the position of the Prisms and Lens to one another must be corrected, and then if by the help of a long Beam of Wood, as is represented in the Figure, or by a Tube, or some other such instrument made for that purpose, they be made fast in that situation, you may try all the same Experiments in this compounded Beam of Light XY, which in the foregoing Experiments have been made in the Sun's direct Light. For this compounded Beam of Light has the same appearance, and is endowed with all the same Properties with a direct Beam of the Sun's Light, so far as my Observation reaches. And in trying Experiments in this Beam you may by stopping any of the Colours p, q, r, s and t, at the Lens, see how the Colours produced in the Experiments are no other than those which the rays had at the Lens before they entered the composition of this Beam: And by consequence that they arise not from any new modifications of the Light by refractions and reflexions, but from the various separations and mixtures of the rays originally endowed with their colour-making qualities.

So, for instance, having with a Lens $4\frac{1}{4}$ Inches broad, and two Prisms on either Hand $6\frac{1}{4}$ Feet distant from the Lens, made such a Beam of compounded Light: to
examin

examin the reason
refracted this com
another Prism HI
matick Colours PC
hind. And then
r, s, t, at the Lens,
vanish at the Paper
the Lens, the purp
and the rest of the
unless perhaps the
it at the Lens might
lowing refractions.
upon the Lens, the
nish, and so of the
the white Beam of
ve Lights variou
lours which afterw
ctions are no other
was compounded.
kh generates the
not by changing th
by separating the r
qualities before the
fracted Beam white
which were of one
ther upon the Paper
So again, to exam
tural Bodies, I plac
XY, and found th
their own Colours
that those Colours
same Colours at th